

Descriptive Set Theory and Its Applications

Alexander S. Kechris

COPENHAGEN; JUNE 2016

Descriptive Set Theory and Its Applications

Over the past 30 years, there has been extensive work on the interactions and applications of descriptive set theory to other areas of mathematics, including analysis, dynamical systems, model theory, topological groups and combinatorics. My goal in these lectures is to give a taste of this area of research.

The first two lectures will concentrate on some of the earlier work in the 1980's and 1990's and will include an extensive historical background. The last two lectures will discuss areas of research that have been intensely studied in the past two decades.

These lectures require minimal background and should be understood by anyone familiar with the basics of descriptive set theory, measure theory and functional analysis. Also the four lectures are essentially independent of each other.

Descriptive Set Theory and Its Applications

Over the past 30 years, there has been extensive work on the interactions and applications of descriptive set theory to other areas of mathematics, including analysis, dynamical systems, model theory, topological groups and combinatorics. My goal in these lectures is to give a taste of this area of research.

The first two lectures will concentrate on some of the earlier work in the 1980's and 1990's and will include an extensive historical background. The last two lectures will discuss areas of research that have been intensely studied in the past two decades.

These lectures require minimal background and should be understood by anyone familiar with the basics of descriptive set theory, measure theory and functional analysis. Also the four lectures are essentially independent of each other.

Descriptive Set Theory and Its Applications

Over the past 30 years, there has been extensive work on the interactions and applications of descriptive set theory to other areas of mathematics, including analysis, dynamical systems, model theory, topological groups and combinatorics. My goal in these lectures is to give a taste of this area of research.

The first two lectures will concentrate on some of the earlier work in the 1980's and 1990's and will include an extensive historical background. The last two lectures will discuss areas of research that have been intensely studied in the past two decades.

These lectures require minimal background and should be understood by anyone familiar with the basics of descriptive set theory, measure theory and functional analysis. Also the four lectures are essentially independent of each other.

Descriptive Set Theory and Its Applications

Here are the topics of the four lectures:

- 1 Set theory and trigonometric series
- 2 On Banach's Problem 50 in the Scottish Book
- 3 The complexity of classification problems in ergodic theory
- 4 Descriptive graph combinatorics

Descriptive Set Theory and Its Applications

Here are the topics of the four lectures:

- 1 **Set theory and trigonometric series**
- 2 On Banach's Problem 50 in the Scottish Book
- 3 The complexity of classification problems in ergodic theory
- 4 Descriptive graph combinatorics

Descriptive Set Theory and Its Applications

Here are the topics of the four lectures:

- 1 **Set theory and trigonometric series**
- 2 **On Banach's Problem 50 in the Scottish Book**
- 3 The complexity of classification problems in ergodic theory
- 4 Descriptive graph combinatorics

Descriptive Set Theory and Its Applications

Here are the topics of the four lectures:

- 1 **Set theory and trigonometric series**
- 2 **On Banach's Problem 50 in the Scottish Book**
- 3 **The complexity of classification problems in ergodic theory**
- 4 **Descriptive graph combinatorics**

Descriptive Set Theory and Its Applications

Here are the topics of the four lectures:

- 1 **Set theory and trigonometric series**
- 2 **On Banach's Problem 50 in the Scottish Book**
- 3 **The complexity of classification problems in ergodic theory**
- 4 **Descriptive graph combinatorics**

What is descriptive set theory?

Descriptive set theory is the study of definable sets and functions in Polish (complete, separable metric) spaces, like the Euclidean spaces, the Baire space, Hilbert space and more generally separable Banach spaces, second countable locally compact groups, etc.

In this theory sets are classified in hierarchies according to the complexity of their definitions and the structure of sets at each level of these hierarchies is systematically studied.

What is descriptive set theory?

Descriptive set theory is the study of definable sets and functions in Polish (complete, separable metric) spaces, like the Euclidean spaces, the Baire space, Hilbert space and more generally separable Banach spaces, second countable locally compact groups, etc.

In this theory sets are classified in hierarchies according to the complexity of their definitions and the structure of sets at each level of these hierarchies is systematically studied.

What is descriptive set theory?

Of particular importance are the Borel and projective sets. The Borel sets are obtained from the open sets by applying repeatedly countable Boolean operations and the projective sets are obtained from the Borel sets by the operations of complementation and projection.

These classes of sets are ramified in natural hierarchies as follows:



$\Sigma_1^1 = \textit{analytic sets}$, $\Pi_1^1 = \textit{co-analytic sets}$

What is descriptive set theory?

Of particular importance are the Borel and projective sets. The Borel sets are obtained from the open sets by applying repeatedly countable Boolean operations and the projective sets are obtained from the Borel sets by the operations of complementation and projection.

These classes of sets are ramified in natural hierarchies as follows:

$$\begin{array}{cccc}
 \Sigma_1^0 & \Sigma_2^0 & \Sigma_3^0 & \dots \\
 \Pi_1^0 & \Pi_2^0 & \Pi_3^0 & \dots \\
 \underbrace{\hspace{10em}} & & & \\
 \text{Borel} & & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 \Sigma_1^1 & \Sigma_2^1 & \dots \\
 \Pi_1^1 & \Pi_2^1 & \dots \\
 \underbrace{\hspace{5em}} & & \\
 \text{Projective} & &
 \end{array}$$

$\Sigma_1^1 = \text{analytic sets}$, $\Pi_1^1 = \text{co-analytic sets}$

What is descriptive set theory?

Of particular importance are the Borel and projective sets. The Borel sets are obtained from the open sets by applying repeatedly countable Boolean operations and the projective sets are obtained from the Borel sets by the operations of complementation and projection.

These classes of sets are ramified in natural hierarchies as follows:

$$\begin{array}{cccc}
 \Sigma_1^0 & \Sigma_2^0 & \Sigma_3^0 & \dots \\
 \Pi_1^0 & \Pi_2^0 & \Pi_3^0 & \dots \\
 \underbrace{\hspace{10em}} & & & \\
 \text{Borel} & & &
 \end{array}
 \qquad
 \begin{array}{ccc}
 \Sigma_1^1 & \Sigma_2^1 & \dots \\
 \Pi_1^1 & \Pi_2^1 & \dots \\
 \underbrace{\hspace{5em}} & & \\
 \text{Projective} & &
 \end{array}$$

$\Sigma_1^1 = \text{analytic sets}$, $\Pi_1^1 = \text{co-analytic sets}$

What is descriptive set theory?

Intuitively, sets whose membership is characterized in “explicit” terms, even allowing countable operations, are Borel.

Here is a classical example of a co-analytic non-Borel set in analysis:

Theorem (Mazurkiewicz, 1936)

In the space $C([0, 1])$, the set of (everywhere) differentiable functions is co-analytic but not Borel.

What is descriptive set theory?

Intuitively, sets whose membership is characterized in “explicit” terms, even allowing countable operations, are Borel.

Here is a classical example of a co-analytic non-Borel set in analysis:

Theorem (Mazurkiewicz, 1936)

In the space $C([0, 1])$, the set of (everywhere) differentiable functions is co-analytic but not Borel.

What is descriptive set theory?

Intuitively, sets whose membership is characterized in “explicit” terms, even allowing countable operations, are Borel.

Here is a classical example of a co-analytic non-Borel set in analysis:

Theorem (Mazurkiewicz, 1936)

In the space $C([0, 1])$, the set of (everywhere) differentiable functions is co-analytic but not Borel.

Lecture I. Set theory and trigonometric series

Trigonometric series

Definition

A **trigonometric series** is an expression of the form

$$s \sim \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos nx + b_n \sin nx$$

or equivalently

$$s \sim \sum_{n=-\infty}^{\infty} c_n e^{inx}, x \in \mathbb{T} = \mathbb{R}/2\pi\mathbb{Z}$$

Trigonometric series

Definition

A **Fourier series** is an expression of the form

$$s \sim \sum_{n=-\infty}^{\infty} \hat{f}(n)e^{inx}, f \in L^1(\mathbb{T}),$$

where $\hat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(x)e^{-inx} dx$.

Definition

A **Fourier-Stieltjes series** is an expression of the form

$$s \sim \sum_{n=-\infty}^{\infty} \hat{\mu}(n)e^{inx}, \mu \in M(\mathbb{T}),$$

where $\hat{\mu}(n) = \int e^{-inx} d\mu$.

Trigonometric series

Definition

A **Fourier series** is an expression of the form

$$s \sim \sum_{n=-\infty}^{\infty} \hat{f}(n)e^{inx}, f \in L^1(\mathbb{T}),$$

where $\hat{f}(n) = \frac{1}{2\pi} \int_0^{2\pi} f(x)e^{-inx} dx$.

Definition

A **Fourier-Stieltjes series** is an expression of the form

$$s \sim \sum_{n=-\infty}^{\infty} \hat{\mu}(n)e^{inx}, \mu \in M(\mathbb{T}),$$

where $\hat{\mu}(n) = \int e^{-inx} d\mu$.

I: Riemann, Heine and Cantor

Riemann, Heine and Cantor



(RIEMANN, 1826–1866)

- Riemann Habilitationsschrift (1854)

Study of the structure of functions that can be represented by trigonometric series

$$f(x) = \sum_{n=-\infty}^{\infty} c_n e^{inx}$$

Riemann, Heine and Cantor

Suggests three general problems:

- (The Uniqueness Problem) Is such an expansion unique?
- (The Characterization Problem) Can one characterize the functions that have a trigonometric expansion?
- (The Coefficient Problem) How does one “compute” the coefficients of the expansion from the function?

Riemann, Heine and Cantor

Suggests three general problems:

- **(The Uniqueness Problem)** Is such an expansion unique?
- **(The Characterization Problem)** Can one characterize the functions that have a trigonometric expansion?
- **(The Coefficient Problem)** How does one “compute” the coefficients of the expansion from the function?

Riemann, Heine and Cantor

Suggests three general problems:

- (The Uniqueness Problem) Is such an expansion unique?
- (The Characterization Problem) Can one characterize the functions that have a trigonometric expansion?
- (The Coefficient Problem) How does one “compute” the coefficients of the expansion from the function?

Riemann, Heine and Cantor

Suggests three general problems:

- (The Uniqueness Problem) Is such an expansion unique?
- (The Characterization Problem) Can one characterize the functions that have a trigonometric expansion?
- (The Coefficient Problem) How does one “compute” the coefficients of the expansion from the function?

Riemann, Heine and Cantor

Will concentrate on the Uniqueness Problem but here are some comments on the other two problems.

- (The Characterization Problem) Even for continuous functions, although there are many well-known sufficient criteria for the expansion in a trigonometric series, one can argue (on the basis of the following result) that no reasonable exact criteria can be found.

Theorem (Ajtai-K, 1987)

In the space $C(\mathbb{T})$, the set of functions that can be expanded in a trigonometric series is co-analytic but not Borel.

Riemann, Heine and Cantor

Will concentrate on the Uniqueness Problem but here are some comments on the other two problems.

- (The Characterization Problem) Even for continuous functions, although there are many well-known sufficient criteria for the expansion in a trigonometric series, one can argue (on the basis of the following result) that no reasonable exact criteria can be found.

Theorem (Ajtai-K, 1987)

In the space $C(\mathbb{T})$, the set of functions that can be expanded in a trigonometric series is co-analytic but not Borel.

Riemann, Heine and Cantor

Will concentrate on the Uniqueness Problem but here are some comments on the other two problems.

- (The Characterization Problem) Even for continuous functions, although there are many well-known sufficient criteria for the expansion in a trigonometric series, one can argue (on the basis of the following result) that no reasonable exact criteria can be found.

Theorem (Ajtai-K, 1987)

In the space $C(\mathbb{T})$, the set of functions that can be expanded in a trigonometric series is co-analytic but not Borel.

Riemann, Heine and Cantor

- (The Coefficient Problem) If an integrable function can be represented by a trigonometric series, then the coefficients are its Fourier coefficients (de la Vallée-Poussin). However there are everywhere convergent series, like

$$\sum_{n=2}^{\infty} \frac{\sin(nx)}{\log n},$$

whose sum is not integrable. Denjoy from 1941 to 1949 wrote a 700 (!) page book describing a general procedure for computing the coefficients.

Riemann, Heine and Cantor



(HEINE, 1821-1881)



(CANTOR, 1845-1918)

Heine suggested to Cantor to study the Uniqueness Problem.

Theorem (Cantor, 1870)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, then $c_n = 0, \forall n$.

Theorem (Cantor, 1872)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, except on a countable closed set of finite Cantor-Bendixson rank, then $c_n = 0, \forall n$.

Riemann, Heine and Cantor



(HEINE, 1821-1881)



(CANTOR, 1845-1918)

Heine suggested to Cantor to study the Uniqueness Problem.

Theorem (Cantor, 1870)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, then $c_n = 0, \forall n$.

Theorem (Cantor, 1872)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, except on a countable closed set of finite Cantor-Bendixson rank, then $c_n = 0, \forall n$.

Riemann, Heine and Cantor



(HEINE, 1821-1881)



(CANTOR, 1845-1918)

Heine suggested to Cantor to study the Uniqueness Problem.

Theorem (Cantor, 1870)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, then $c_n = 0, \forall n$.

Theorem (Cantor, 1872)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, except on a countable closed set of finite Cantor-Bendixson rank, then $c_n = 0, \forall n$.

Riemann, Heine and Cantor



(LEBESGUE, 1875-1941)



(W.H. YOUNG, 1863-1942)

Theorem (Lebesgue, 1903)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, except on a closed countable set, then $c_n = 0, \forall n$.

Theorem (W.H.Young, 1909)

If $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \in \mathbb{T}$, except on a countable set, then $c_n = 0, \forall n$.

Sets of uniqueness

Definition

A (Borel) set $A \subseteq \mathbb{T}$ is called a **set of uniqueness** if $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \notin A$, implies $c_n = 0, \forall n$. Otherwise it is called a **set of multiplicity**.

We denote by \mathcal{U} the class of sets of uniqueness and by \mathcal{M} the class of sets of multiplicity. Thus

$$\text{countable} \subseteq \mathcal{U} \subseteq (\text{Lebesgue}) \text{ null.}$$

- Is $\mathcal{U} = \text{countable}$?
- Is $\mathcal{U} = (\text{Lebesgue}) \text{ null}$?

Sets of uniqueness

Definition

A (Borel) set $A \subseteq \mathbb{T}$ is called a **set of uniqueness** if $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \notin A$, implies $c_n = 0, \forall n$. Otherwise it is called a **set of multiplicity**.

We denote by \mathcal{U} the class of sets of uniqueness and by \mathcal{M} the class of sets of multiplicity. Thus

$$\text{countable} \subseteq \mathcal{U} \subseteq (\text{Lebesgue}) \text{ null.}$$

- Is $\mathcal{U} = \text{countable}$?
- Is $\mathcal{U} = (\text{Lebesgue}) \text{ null}$?

Sets of uniqueness

Definition

A (Borel) set $A \subseteq \mathbb{T}$ is called a **set of uniqueness** if $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \notin A$, implies $c_n = 0, \forall n$. Otherwise it is called a **set of multiplicity**.

We denote by \mathcal{U} the class of sets of uniqueness and by \mathcal{M} the class of sets of multiplicity. Thus

$$\text{countable} \subseteq \mathcal{U} \subseteq (\text{Lebesgue}) \text{ null.}$$

- Is $\mathcal{U} = \text{countable}$?
- Is $\mathcal{U} = (\text{Lebesgue}) \text{ null}$?

Sets of uniqueness

Definition

A (Borel) set $A \subseteq \mathbb{T}$ is called a **set of uniqueness** if $\sum_{n=-\infty}^{\infty} c_n e^{inx} = 0, \forall x \notin A$, implies $c_n = 0, \forall n$. Otherwise it is called a **set of multiplicity**.

We denote by \mathcal{U} the class of sets of uniqueness and by \mathcal{M} the class of sets of multiplicity. Thus

$$\text{countable} \subseteq \mathcal{U} \subseteq (\text{Lebesgue}) \text{ null.}$$

- Is $\mathcal{U} = \text{countable}$?
- Is $\mathcal{U} = (\text{Lebesgue}) \text{ null}$?

II. The Russian and Polish Schools (mid 1910's - mid 1930's)

The Russian and Polish Schools

The structure of sets of uniqueness was investigated intensely during that period by the Russian school of Luzin, Menshov and Bari, and the Polish school of Rajchman, Zygmund and Marcinkiewicz.

The Russian and Polish Schools



(LUZIN, 1883-1950)



(MENSHOV, 1892-1988)



(BARI, 1901-1961)

The Russian and Polish Schools



(RAJCHMAN, 1890-1940)



(ZYGmund, 1900-1992)



(MARCINKIEWICZ, 1910-1940)

The Russian and Polish Schools

Theorem (Bari, Zygmund 1923)

The union of countably many closed sets of uniqueness is also a set of uniqueness.

The Russian and Polish Schools

Definition

Given real numbers ξ_1, ξ_2, \dots , with $0 < \xi_n < 1/2$, denote by $E_{\xi_1, \xi_2, \dots}$ the Cantor-type set (in \mathbb{T}) constructed with successive ratios of dissection ξ_1, ξ_2, \dots . We also let $E_\xi = E_{\xi, \xi, \dots}$. In particular, $E_{1/3}$ is the usual Cantor set.

The Russian and Polish Schools

Theorem (Menshov, 1916)

There is a closed null set of multiplicity. In fact, $E_{\xi_1, \xi_2, \dots}$, with $\xi_n = \frac{(n+1)}{2(n+2)}$, is such a set.

Theorem (Bari, Rajchman, 1921-1923)

There are perfect sets of uniqueness. In fact, $E_{1/3}$ is such a set.

It follows that

$$\text{countable } \not\subseteq \mathcal{U} \not\subseteq \text{null.}$$

Thus by the 1920's it had become clear that the class of (even closed) sets of uniqueness is hard to delineate in terms of classical notions of smallness in analysis.

The Russian and Polish Schools

Theorem (Menshov, 1916)

There is a closed null set of multiplicity. In fact, $E_{\xi_1, \xi_2, \dots}$, with $\xi_n = \frac{(n+1)}{2(n+2)}$, is such a set.

Theorem (Bari, Rajchman, 1921-1923)

There are perfect sets of uniqueness. In fact, $E_{1/3}$ is such a set.

It follows that

$$\text{countable } \not\subseteq \mathcal{U} \not\subseteq \text{null.}$$

Thus by the 1920's it had become clear that the class of (even closed) sets of uniqueness is hard to delineate in terms of classical notions of smallness in analysis.

The Russian and Polish Schools

Theorem (Menshov, 1916)

There is a closed null set of multiplicity. In fact, $E_{\xi_1, \xi_2, \dots}$, with $\xi_n = \frac{(n+1)}{2(n+2)}$, is such a set.

Theorem (Bari, Rajchman, 1921-1923)

There are perfect sets of uniqueness. In fact, $E_{1/3}$ is such a set.

It follows that

$$\text{countable } \not\subseteq \mathcal{U} \not\subseteq \text{null.}$$

Thus by the 1920's it had become clear that the class of (even closed) sets of uniqueness is hard to delineate in terms of classical notions of smallness in analysis.

The Russian and Polish Schools

Bari's memoir in 1927 stated some classical problems on sets of uniqueness.

- **(The Characterization Problem)** Find necessary and sufficient conditions for a perfect set to be a set of uniqueness.

It appears that the intended meaning was to ask for geometric, analytic (or, as we will see later, even number theoretic) structural properties of a given perfect set, expressed “explicitly” in terms of some standard description of it, that will determine whether it is a set of uniqueness or multiplicity.

The Russian and Polish Schools

Bari's memoir in 1927 stated some classical problems on sets of uniqueness.

- **(The Characterization Problem) Find necessary and sufficient conditions for a perfect set to be a set of uniqueness.**

It appears that the intended meaning was to ask for geometric, analytic (or, as we will see later, even number theoretic) structural properties of a given perfect set, expressed “explicitly” in terms of some standard description of it, that will determine whether it is a set of uniqueness or multiplicity.

The Russian and Polish Schools

- **(The Union Problem) Is the union of two (Borel) sets of uniqueness also a set of uniqueness?**

The first open case is that of two G_δ sets.

- **(The Category Problem) Is every (Borel) set of uniqueness of the first category (meager)?**

The Russian and Polish Schools

- **(The Union Problem)** Is the union of two (Borel) sets of uniqueness also a set of uniqueness?

The first open case is that of two G_δ sets.

- **(The Category Problem)** Is every (Borel) set of uniqueness of the first category (meager)?

III. Thin sets in harmonic analysis (early 1950's - mid 1970's)

Thin sets in harmonic analysis

During that period there was an explosion of research into the structure of thin sets in harmonic analysis, including the study of closed sets of uniqueness.

Piatetski-Shapiro in 1952-54 introduced functional analysis methods into the subject of uniqueness. This has become the standard language of the subject since then.



(PIATETSKI-SHAPIRO, 1929-2009)

Thin sets in harmonic analysis

During that period there was an explosion of research into the structure of thin sets in harmonic analysis, including the study of closed sets of uniqueness.

Piatetski-Shapiro in 1952-54 introduced functional analysis methods into the subject of uniqueness. This has become the standard language of the subject since then.



(PIATETSKI-SHAPIRO, 1929-2009)

Thin sets in harmonic analysis

Definition

We denote by $A(\mathbb{T})$ the Banach algebra of continuous functions with absolutely convergent Fourier series. This is of course the same as $\ell^1(\mathbb{Z})$. Its dual is the space $\ell^\infty(\mathbb{Z})$, which in this context is called the space of **pseudomeasures** and denoted by **PM**. Its predual is the space $c_0(\mathbb{Z})$, which in this context is called the space of **pseudofunctions** and denoted by **PF**.

Definition

The **support** of a pseudomeasure S is the complement of the largest open set on which S vanishes, i.e., annihilates all functions in $A(\mathbb{T})$ supported by it.

Thin sets in harmonic analysis

Definition

We denote by $A(\mathbb{T})$ the Banach algebra of continuous functions with absolutely convergent Fourier series. This is of course the same as $\ell^1(\mathbb{Z})$. Its dual is the space $\ell^\infty(\mathbb{Z})$, which in this context is called the space of **pseudomeasures** and denoted by **PM**. Its predual is the space $c_0(\mathbb{Z})$, which in this context is called the space of **pseudofunctions** and denoted by **PF**.

Definition

The **support** of a pseudomeasure S is the complement of the largest open set on which S vanishes, i.e., annihilates all functions in $A(\mathbb{T})$ supported by it.

Thin sets in harmonic analysis

Piatetski-Shapiro's reformulation of the concept of closed set of uniqueness.

Theorem (Piatetski-Shapiro, 1952)

A closed set E is a set of uniqueness iff it does not support a non-0 pseudofunction.

Thin sets in harmonic analysis

At this point it is time to introduce an important variation of the concept of set of uniqueness, which really goes back to Menshov's work. His example of a null closed set of multiplicity was witnessed by the Fourier-Stieltjes series of a (probability Borel) measure. Such a set is called a set of strict multiplicity.

Definition

A (Borel) set is called a set of **extended uniqueness** if it satisfies uniqueness for Fourier-Stieltjes series of measures. Otherwise it is called a set of **strict multiplicity**. The class of sets of extended uniqueness is denoted by \mathcal{U}_0 and the class of sets of strict multiplicity is denoted by \mathcal{M}_0 .

Thin sets in harmonic analysis

At this point it is time to introduce an important variation of the concept of set of uniqueness, which really goes back to Menshov's work. His example of a null closed set of multiplicity was witnessed by the Fourier-Stieltjes series of a (probability Borel) measure. Such a set is called a set of strict multiplicity.

Definition

A (Borel) set is called a set of **extended uniqueness** if it satisfies uniqueness for Fourier-Stieltjes series of measures. Otherwise it is called a set of **strict multiplicity**. The class of sets of extended uniqueness is denoted by \mathcal{U}_0 and the class of sets of strict multiplicity is denoted by \mathcal{M}_0 .

Thin sets in harmonic analysis

Definition

A **Rajchman measure** is a measure whose Fourier-Stieltjes coefficients converge to 0, i.e., form a pseudofunction.

Lebesgue measure is of this form and a Rajchman measure is thought of as a measure with “large support”. However, Menshov showed that there are singular Rajchman measures. In terms of Rajchman measures, the sets of extended uniqueness are exactly those that are null for all such measures.

Theorem (Piatetski-Shapiro, 1954)

There is a closed set of extended uniqueness but not of uniqueness.

Thin sets in harmonic analysis

Definition

A **Rajchman measure** is a measure whose Fourier-Stieltjes coefficients converge to 0, i.e., form a pseudofunction.

Lebesgue measure is of this form and a Rajchman measure is thought of as a measure with “large support”. However, Menshov showed that there are singular Rajchman measures. In terms of Rajchman measures, the sets of extended uniqueness are exactly those that are null for all such measures.

Theorem (Piatetski-Shapiro, 1954)

There is a closed set of extended uniqueness but not of uniqueness.

Thin sets in harmonic analysis

Definition

A **Rajchman measure** is a measure whose Fourier-Stieltjes coefficients converge to 0, i.e., form a pseudofunction.

Lebesgue measure is of this form and a Rajchman measure is thought of as a measure with “large support”. However, Menshov showed that there are singular Rajchman measures. In terms of Rajchman measures, the sets of extended uniqueness are exactly those that are null for all such measures.

Theorem (Piatetski-Shapiro, 1954)

There is a closed set of extended uniqueness but not of uniqueness.

Thin sets in harmonic analysis

This result of Piatetski-Shapiro was amplified in the work of Körner in the early 1970's, who solved a major problem at that time by constructing a particular kind of closed thin set, called a *Helson set*, which is of multiplicity. As Helson, 1954 had already shown that these sets are of extended uniqueness, this also implied the Piatetski-Shapiro theorem. This result of Körner was one of the last major results of that period. Its original proof was extremely complicated and despite a major simplification by Kaufman it remains a subtle and difficult result.

Thin sets in harmonic analysis

During that period there was also a major advance in the characterization problem.

Definition

A real number is called a **Pisot** (or **Pisot-Vijayaraghavan**) number if it is an algebraic integer > 1 all of whose conjugates have absolute value < 1 .

Examples: The integers > 1 and the golden mean.

Intuitively, these are numbers whose powers approach integers.

Thin sets in harmonic analysis

During that period there was also a major advance in the characterization problem.

Definition

A real number is called a **Pisot** (or **Pisot-Vijayaraghavan**) number if it is an algebraic integer > 1 all of whose conjugates have absolute value < 1 .

Examples: The integers > 1 and the golden mean.

Intuitively, these are numbers whose powers approach integers.

Thin sets in harmonic analysis

During that period there was also a major advance in the characterization problem.

Definition

A real number is called a **Pisot** (or **Pisot-Vijayaraghavan**) number if it is an algebraic integer > 1 all of whose conjugates have absolute value < 1 .

Examples: The integers > 1 and the golden mean.

Intuitively, these are numbers whose powers approach integers.

Thin sets in harmonic analysis



(SALEM, 1898-1963)

Theorem (Salem, 1944)

The Pisot numbers form a closed set of algebraic integers.

Theorem (Salem-Zygmund, 1955)

The set E_ξ is a set of uniqueness iff $1/\xi$ is a Pisot number.

Thin sets in harmonic analysis



(SALEM, 1898-1963)

Theorem (Salem, 1944)

The Pisot numbers form a closed set of algebraic integers.

Theorem (Salem-Zygmund, 1955)

The set E_ξ is a set of uniqueness iff $1/\xi$ is a Pisot number.

IV. Applications of descriptive set theory (mid 1980's - mid 1990's)

Applications of descriptive set theory

We have seen that the problems of uniqueness have involved ideas from many subjects, such as classical real analysis, modern harmonic analysis, functional analysis, number theory, etc. Although set theory owes its origin to Cantor's work on the uniqueness problem, relatively little contact existed between set theory and the study of sets of uniqueness until the 1980's, when ideas from descriptive set theory, were brought to bear in the study of this subject. This is interesting since descriptive set theory was originally developed in the same Russian and Polish schools during the same period 1915-1935.

Applications of descriptive set theory

Luzin's school was concerned with what was then called the *theory of real functions* and there was at that time a distinction between the so-called *metric theory* (differentiation, integration, trigonometric series, etc.) and the *descriptive theory* (called today descriptive set theory). Strangely enough, according to Kolmogorov, who was a member of that school, Luzin divided his students to those that would study the metric theory and those that would study the descriptive one. (Kolmogorov actually violated this rule and worked on both.) In the following years the subjects drifted apart, the first one practiced by analysts and the second one by logicians. They were brought back together in the 1980's in the study of sets of uniqueness.

Applications of descriptive set theory

In the 1980's and 1990's methods of descriptive set theory were combined with previous work in analysis to the study of sets of uniqueness. This was primarily developed in series of papers (and unpublished work) of the following mathematicians: Debs-Saint Raymond, Kaufman, K-Louveau, K-Louveau-Woodin, Solovay.

The main point is that descriptive set theory allows one to develop a “global” theory of closed sets of uniqueness, with many applications to the classical theory.

Applications of descriptive set theory

In the 1980's and 1990's methods of descriptive set theory were combined with previous work in analysis to the study of sets of uniqueness. This was primarily developed in series of papers (and unpublished work) of the following mathematicians: Debs-Saint Raymond, Kaufman, K-Louveau, K-Louveau-Woodin, Solovay.

The main point is that descriptive set theory allows one to develop a “global” theory of closed sets of uniqueness, with many applications to the classical theory.

Applications of descriptive set theory

The appropriate space here is the compact metric space $K(\mathbb{T})$ of closed subsets of the circle, with the usual Hausdorff metric. One studies the structure of the following two subsets of this space:

- $U = \{E \in K(\mathbb{T}) : E \text{ is a set of uniqueness}\}$
- $U_0 = \{E \in K(\mathbb{T}) : E \text{ is a set of extended uniqueness}\}$

The global theory can be encapsulated in the following main theorem, whose proof is contained in a series of papers of the above authors. It states three basic principles that describe the structure of the classes U , U_0 .

Applications of descriptive set theory

The appropriate space here is the compact metric space $K(\mathbb{T})$ of closed subsets of the circle, with the usual Hausdorff metric. One studies the structure of the following two subsets of this space:

- $U = \{E \in K(\mathbb{T}) : E \text{ is a set of uniqueness}\}$
- $U_0 = \{E \in K(\mathbb{T}) : E \text{ is a set of extended uniqueness}\}$


The global theory can be encapsulated in the following main theorem, whose proof is contained in a series of papers of the above authors. It states three basic principles that describe the structure of the classes U , U_0 .

Applications of descriptive set theory

Theorem

- a) (Stability property) *The sets U, U_0 are calibrated σ -ideals.*
- b) (Descriptive complexity, I) *Both U, U_0 are co-analytic and locally non-Borel, i.e., for every closed set E not in U (resp., not in U_0) the set $U \cap K(E)$ (resp., $U_0 \cap K(E)$) is not Borel.*
- c) (Descriptive complexity, II) *The σ -ideal U_0 admits a Borel basis but the σ -ideal U does not.*

Applications of descriptive set theory



Applications of descriptive set theory

This theory has numerous applications both in the solution of classical problems and also in understanding and proving in a new way previously established results.

Applications of descriptive set theory

- (Characterization Problem) One can argue that this has a negative solution in a strong sense, since not only there is no “explicit” characterization of perfect sets of uniqueness of the sought after type but also there is no way to characterize such sets in terms of decomposing them into a countable number of explicitly characterizable components.

Applications of descriptive set theory

- Every known until the early 1980's closed set of uniqueness could be written as a union of a countable sequence of simpler uniqueness sets, belonging to a class denoted by U' . This is a Borel class, so the non-basis theorem shows that there are many new kinds of U -sets. This result can therefore be viewed as a powerful new existence theorem. For example, it answers a question of Piatetski-Shapiro, on the existence of U -sets not expressible as countable unions of so-called $H^{(n)}$ -sets (with varying n).

Applications of descriptive set theory

- (Category Problem) This is completely solved affirmatively in a strong sense, as it follows that every Borel set of *extended uniqueness* is of the first category. Equivalently this means that every Borel set of the second category supports a Rajchman measure. This in turn has several applications, including in particular a unified, new and much simpler way of proving some well-known results in the theory, originally established by various techniques and constructions.

Applications of descriptive set theory

- Menshov's Theorem says that there are (Lebesgue) null sets that support Rajchman measures. This is now clear as it well-known that there are comeager null sets. Thus Menshov's result is seen as a consequence of the orthogonality between measure and category.
- Ivashev-Musatov and Kaufman have extended Menshov's Theorem to show that for any function h there are h -Hausdorff measure 0 closed sets that support Rajchman measures. The same argument as above applies.
- The Kahane-Salem (1964) problem asks whether the set of non-normal numbers supports a Rajchman measure. This was solved affirmatively by Lyons (1986). Again this follows from the fact that the set of non-normal numbers is comeager.

Applications of descriptive set theory

- Menshov's Theorem says that there are (Lebesgue) null sets that support Rajchman measures. This is now clear as it well-known that there are comeager null sets. Thus Menshov's result is seen as a consequence of the orthogonality between measure and category.
- Ivashev-Musatov and Kaufman have extended Menshov's Theorem to show that for any function h there are h -Hausdorff measure 0 closed sets that support Rajchman measures. The same argument as above applies.
- The Kahane-Salem (1964) problem asks whether the set of non-normal numbers supports a Rajchman measure. This was solved affirmatively by Lyons (1986). Again this follows from the fact that the set of non-normal numbers is comeager.

Applications of descriptive set theory

- Menshov's Theorem says that there are (Lebesgue) null sets that support Rajchman measures. This is now clear as it well-known that there are comeager null sets. Thus Menshov's result is seen as a consequence of the orthogonality between measure and category.
- Ivashev-Musatov and Kaufman have extended Menshov's Theorem to show that for any function h there are h -Hausdorff measure 0 closed sets that support Rajchman measures. The same argument as above applies.
- The Kahane-Salem (1964) problem asks whether the set of non-normal numbers supports a Rajchman measure. This was solved affirmatively by Lyons (1986). Again this follows from the fact that the set of non-normal numbers is comeager.

Applications of descriptive set theory

- (The Union Problem) This is still open, even for the union of two G_δ sets. It is mostly believed that there is a counterexample. The preceding theory however implies that from a counterexample one obtains a closed set with properties similar to those obtained by Körner (Helson sets of multiplicity). Thus conceivably Körner's Theorem could be useful in the construction of such a counterexample.

Applications of descriptive set theory

There are several further applications of descriptive set theoretic methods also to other aspects of the subject, e.g., Lyons' characterization of Rajchman measures by their null sets is seen as following from a general descriptive set theoretic result of Mokobodzki about analytic classes of measures. Also such ideas have been applied by S. Kahane (a cousin of J.-P. Kahane) to the solution of some old problems about other types of thin sets in harmonic analysis.

Conclusion

This is where we stand now. Despite the progress made over the last 150 years many mysteries remain. Here are for example some intriguing problems that are still open:

- Where is the dividing line in the Characterization Problem?

E_ξ : characterizable

$E_{\xi_1, \xi_2, \dots}$: ???

E in general: uncharacterizable

- The Union Problem for G_δ sets and for arbitrary Borel sets.
- (The Interior Problem, Bari 1927) Is the concept of set of uniqueness determined by the closed sets, i.e., does a Borel set of multiplicity contain a closed set of multiplicity?

Conclusion

This is where we stand now. Despite the progress made over the last 150 years many mysteries remain. Here are for example some intriguing problems that are still open:

- Where is the dividing line in the Characterization Problem?

E_ξ : characterizable

$E_{\xi_1, \xi_2, \dots}$: ???

E in general: uncharacterizable

- The Union Problem for G_δ sets and for arbitrary Borel sets.
- (The Interior Problem, Bari 1927) Is the concept of set of uniqueness determined by the closed sets, i.e., does a Borel set of multiplicity contain a closed set of multiplicity?

Conclusion

This is where we stand now. Despite the progress made over the last 150 years many mysteries remain. Here are for example some intriguing problems that are still open:

- Where is the dividing line in the Characterization Problem?

E_ξ : characterizable

$E_{\xi_1, \xi_2, \dots}$: ???

E in general: uncharacterizable

- The Union Problem for G_δ sets and for arbitrary Borel sets.
- (The Interior Problem, Bari 1927) Is the concept of set of uniqueness determined by the closed sets, i.e., does a Borel set of multiplicity contain a closed set of multiplicity?

Conclusion

This is where we stand now. Despite the progress made over the last 150 years many mysteries remain. Here are for example some intriguing problems that are still open:

- Where is the dividing line in the Characterization Problem?

E_ξ : characterizable

$E_{\xi_1, \xi_2, \dots}$: ???

E in general: uncharacterizable

- The Union Problem for G_δ sets and for arbitrary Borel sets.
- (The Interior Problem, Bari 1927) Is the concept of set of uniqueness determined by the closed sets, i.e., does a Borel set of multiplicity contain a closed set of multiplicity?

Conclusion

This is where we stand now. Despite the progress made over the last 150 years many mysteries remain. Here are for example some intriguing problems that are still open:

- Where is the dividing line in the Characterization Problem?

E_ξ : characterizable

$E_{\xi_1, \xi_2, \dots}$: ???

E in general: uncharacterizable

- The Union Problem for G_δ sets and for arbitrary Borel sets.
- (The Interior Problem, Bari 1927) Is the concept of set of uniqueness determined by the closed sets, i.e., does a Borel set of multiplicity contain a closed set of multiplicity?